



Han G. Sohn
MD, FACS

Post-Operative Care and Instructions for PE Tubes

1. **DRY EAR PRECAUTIONS** – Do not get water into ears – especially bath water. This can cause an ear infection. Typical showering is OK.
2. Use swim molds to protect ears from water entry if planning on water activities such as swimming/diving.
3. Use prescribed ear drops: Floxin / Ciprodex – 5 drops to each ear (with PE Tube) every 12 hours as directed. When applying drops, remember to pull the ear back and rub the front of the ear to help the drops get into the ear.
4. Drainage and/or bloody discharge from the ears can be expected after surgery. Even months after surgery, bloody discharge can occur and usually indicates that an ear infection is coming on. Please notify our office so that you can be instructed on how to manage this and to schedule an appointment to have your ears checked. Do not use any over the counter eardrops or previously prescribed eardrops, as they may not be safe now that the eardrum is opened with a tube.
5. For one (1) week after surgery: a. Do not blow nose after PE Tube placement. This could cause the tube to be displaced. b. If you must sneeze, try to open your mouth to avoid pushing the PE Tube out accidentally.

We look forward to seeing you on your post-operative appointment ~2 weeks after surgery

410.889.0795

totalent@totalentcare.com

www.totalentcare.com

3333 N. Calvert Street, Suite 680
Baltimore, MD 21218

2227 Old Emmorton Road, Suite 114
Towson, MD 21286

2227 Old Emmorton Road, Suite 114
Bel Air, MD 21015