



Han G. Sohn

MD, FACS

Post-Operative Care and Instructions for Tonsillectomy

For the first 7 to 10 days following surgery, your activity and diet will be somewhat restricted.

ACTIVITY:

1. Remain quiet during the first week, preferably indoors. You can read, play games, and watch television.
2. Do not engage in strenuous activity during the first 10 days following surgery. No running, biking, swimming, etc.
3. Avoid contact with people with colds, fever, flu, sore throats, virus or any other contagious disease. It would delay your recovery to develop an illness during this time.
4. **DO NOT** take any products containing aspirin as this could affect blood clotting. We recommend using Tylenol instead.

DIET:

Remember that this is just a guideline of suggested foods to prepare. It is fine to substitute foods that you prefer. Some patients advance their diet at a quicker rate than others do. All foods should be lukewarm or cool, not hot. Avoid spicy food and acidic types of fruit juices. It is okay to have apple or grape juices.

1. First day: Iced tea, milk, water, soft drinks, ice cream, Jell-O.
2. Second day: Soft foods like custards, pudding, and cream of wheat, soup broth.
3. Third day: Mashed potatoes, warm cream soups, and poached eggs.
4. Fourth/Fifth day: Soft cereal, soft eggs, and spaghetti.
5. Sixth day and on: Try out more substantial foods that you prefer.

Allow for plenty of fluids and smaller bites to aid in swallowing. **NO** hard foods (peanuts, chips, popcorn etc.) for 3 weeks following your surgery.

410.889.0795

totalent@totalentcare.com

www.totalentcare.com

3333 N. Calvert Street, Suite 680
Baltimore, MD 21218

2227 Old Emmorton Road, Suite 114
Towson, MD 21286

2227 Old Emmorton Road, Suite 114
Bel Air, MD 21015



Han G. Sohn
MD, FACS

WHAT TO EXPECT:

1. **SORE THROAT** - This is usually worse in the morning and at bedtime. It is normally worse on days 5-7 after the operation.
2. **FEVER** - This can be present for the first 2-3 days. It is helpful to encourage a good fluid intake. You can also use Tylenol or Advil for this. If the fever rises over 102° oral temperature, notify us, as we may need to prescribe an antibiotic.
3. **EARACHE**: This is normal even if ear surgery is not performed. The same nerve to your tonsils gives feeling to your ear, as well. The ear feels referred pain or “sympathy” pain.
4. **HIGH-PITCHED (or nasal voice)**: This is common during the first 2-3 weeks following surgery.
5. **BAD BREATH**: This is due to the presence of a membrane or scab where the tonsil was removed.
6. **DIFFICULTY** with swallowing, drooling, and spitting small amounts of bloody mucous can occur in the first few days.
7. **CONSTIPATION** – The narcotic pain medications may cause constipation. Please be aware of this and consult your local pharmacist for an over the counter remedy as needed.

Bleeding can occur about the 7th-10th day after surgery. Report any bleeding immediately so we can discuss treatment options. Ice water gargles and ice pack to neck can sometimes control bleeding. If bleeding is profuse, go to the nearest emergency room. Please call the office for any questions or concerns. We are always available to help you to a speedy recovery.

410.889.0795

totalent@totalentcare.com

www.totalentcare.com

3333 N. Calvert Street, Suite 680
Baltimore, MD 21218

2227 Old Emmorton Road, Suite 114
Towson, MD 21286

2227 Old Emmorton Road, Suite 114
Bel Air, MD 21015